

WHAT ARE NITAZENES?

- → Labeled as the "Frankenstein Opioid" due to its strong effects.
- → Street name ISO, slang names oni
- → Nitazenes are potent synthetic opioids from a drug class known as benzimidazoleopioids.
- → Some nitzenes are estimated to be several times more potent than fentanyl.
- They were developed 60 years ago as potential pain relief medication, but were never approved for clinical use.
- → They were first identified in the unregulated drug supply in America in 2019.

Nitazenes are very strong drugs that may cause overdose even if just a small amount is smoked, snorted, injected, or taken orally.

EXAMPLES OF NITAZENES:

- Etodesnitazene
- Etonitazene
- Etonitazepyne
- Flunitazene
- Isotonitazene
- Metonitazene
- Protonitazene
- 5-Aminoisotonitazene



According to the DEA, two milligrams of fentanyl can constitute a lethal dose. ISO's toxicity can be considerably higher.

WHY THEY ARE A CONCERN

- → The presence of nitazenes in the unregulated drug supply is rising. The frequency with which they are detected is increasing, but there is also an increasing number of nitazene analogs.
- → They tend to be used unintentionally and in unknown quantities because the contents of drugs from the unregulated supply are unpredictable.
- → One of the biggest dangers with these drugs is that they may not be sold as a nitazene. They typically appear unexpectedly in drugs expected to contain other, more traditional opioids (e.g., fentanyl, oxycodone tablets or "down"), often alongside non-medical benzodiazepines (NMBs) as well as illicitly-sold marijuana.
- → Nitazenes can increase the risk of accidental overdose, especially when combined with other substances that suppress breathing and heart rate such as other opioids or benzodiazepines.
- Overdoses involving nitazenes may be difficult to reverse, potentially requiring additional doses of naloxone, but protocols around this are not yet clear.
- Fentanyl test strips cannot detect nitazenes, and detection of nitazenes by pointof-service drug checking requires sensitive equipment that is not always available.
- → Nitazenes are not routinely tested for in post-mortem or urine toxicology, so their relationship to health risks and overdose deaths is difficult to monitor.
- → Naloxone is a key harm reduction tool for reversing overdoses involving opioids. However, it is unclear what the appropriate naloxone response is to reverse an overdose involving a nitazene.

Benzimidazole Opioids	Compared to fentanyl	Compared to morphine
Etonitazene	50x greater	1000x greater
Protonitazene	1.7 to 1.29x greater	130x greater
Isotonitazene	Roughly equal	2.5x greater









